



# Caatinga Ethnoherpetology: Relationships between herpetofauna and people in a semiarid region of northeastern Brazil

<sup>1,2</sup>Lívia Emanuelle Tavares Mendonça, <sup>2</sup>Washington Luiz Silva Vieira, and  
<sup>3,\*</sup>Rômulo Romeu Nóbrega Alves

<sup>1,3</sup>Departamento de Biologia, Universidade Estadual da Paraíba, Av. das Baraúnas, 351/Campus Universitário Bodocongó, 58109-753 Campina Grande, PB, BRAZIL <sup>1,2</sup> Departamento de Sistemática e Ecologia da Universidade Federal da Paraíba, Programa de Pós-Graduação em Ciências Biológicas (Zoologia), Laboratório de Ecofisiologia Animal, 58051-900 João Pessoa, PB, BRAZIL

**Abstract.**—We investigated the interactions between humans and herpetofauna in the semiarid region of Paraíba State, Brazil. Data were obtained by means of interviews with 124 hunters or ex-hunters using semi-structured questionnaires, complemented by informal conversations. We recorded 18 species (17 reptiles and one amphibian) that local human populations interact with because they have some utilitarian value or because of conflicting relations with local inhabitants. Implementation of conservation measures aimed at the herpetofauna in this region is particularly difficult due to the aversion that local people hold toward many of these species. Therefore, environmental education strategies should be adopted. These efforts should not be solely directed at species subject to hunting, but should be all-inclusive and take into consideration the cultural, social, and utilitarian role that governs the interactions of human populations and the herpetofauna of the Caatinga.

**Key words.** Caatinga, conservation, ethnobiology, ethnozoology, hunting, reptiles, wildlife use

Citation: Mendonça LET, Vieira WLS, Alves RRN. 2014. Caatinga Ethnoherpetology: Relationships between herpetofauna and people in a semiarid region of northeastern Brazil. *Amphibian & Reptile Conservation* 8(1) [General Section]: 24–32 (e78).

## Introduction

Humans and herpetofauna (amphibians and reptiles) have interacted for millennia, virtually wherever they have been in contact (Alves et al. 2013b). As a result, interactions between humans and these animals are quite varied, encompassing utilitarian, symbolic, and conflicting aspects (Alves et al. 2008, 2009a, 2012b, c; Fernandes-Ferreira et al. 2012a; Franke and Telecky 2001; Klemens and Thorbjarnarson 1995; Morris and Morris 1965; Moura et al. 2010; Schlaepfer et al. 2005). Such interactions can be studied through ethnoherpetology, a subdivision of ethnozoology, which examines the relationships between human cultures and herpetofauna (Bertrand 1997; Das 1998; Goodman and Hobbs 1994; Speck 1946). Ethnozoological studies can aid in the evaluation of the impacts human populations have on native animal species and in the development of sustainable management plans, and thus, they are essential to conservation efforts (Alves 2012; Alves and Souto 2011).

Caatinga is the name given to the semiarid region that occupies the largest portion of Northeast Brazil and represents one of the major examples of a semiarid environment in the Neotropical region (Albuquerque et al. 2012; Alves et al. 2012b). In this biome, 205 herpetofaunal species have been recorded (65 amphibians, 66 lizards,

12 amphisbaenids, 53 snakes, five testudines, and four crocodylians), many of which interact with local human populations, where they furnish products exploited by the local people or are hunted and killed due to conflicting relations with people (Alves et al. 2009b, 2012a, b, c; Barbosa et al. 2011; Fernandes-Ferreira et al. 2013). In this context, understanding of the relations between humans and the herpetofauna of the region is an important step in designing strategies for management and sustainable use, and should consider the ecological, economic, and cultural aspects associated with these interactions.

Ethnoherpetological studies have only recently begun in Caatinga, although general ethnozoological research indicates that reptiles and amphibians are hunted by rural and urban populations of the region (Albuquerque et al. 2012; Alves et al. 2012b; Fernandes-Ferreira et al. 2012a). In an effort to contribute to our ethnoherpetological knowledge and its implications in the semiarid region of northeastern Brazil, we investigated the interactions between humans and herpetofauna in the municipality of Pocinhos in the semi-arid region of Paraíba State (PB). Our aim was to record the patterns of interactions of the local people with representatives of this animal group in the region. This information may be used to enhance conservation of the Caatinga's herpetofauna.

**Correspondence.** Email: [\\*romulo\\_nobrega@yahoo.com.br](mailto:*romulo_nobrega@yahoo.com.br)

## Materials and Methods

### Study area

The present study was carried out in the municipality of Pocinhos, located in the semi-arid region of Paraíba State, Brazil (Fig. 1; Ribeiro 2003). Pocinhos is 630 km<sup>2</sup> in area, with approximately 17,032 inhabitants. Average annual temperature is 23 °C, which varies little throughout the year. The region has a very low rainfall rate, fluctuating annually between 400 and 600 mm. The climate is hot, semi-arid, with rainfall in the autumn and winter months (Ribeiro 2003) and the vegetation is dominated by sub-deciduous and deciduous forests typical of semi-arid regions (Alves et al. 2009b; Ribeiro 2003).

### Procedures

The study was conducted in the period of June 2010 to June 2011. The information was obtained by means of interviews with hunters or ex-hunters using semistructured questionnaires, complemented by informal conversations (Bernard 1994; Huntington 2000). The selection of informants was done by the “snowball” sampling technique (Bailey 1994), where from the initial contact, an informant indicates another who in turn indicates still another and so forth. Before each interview, the nature and objectives of the research were explained, and the interviewees gave their permission to record the information, by signing an informed consent form.

The questionnaires were applied to 124 hunters from the municipality, of which 98 (79%) live in urban areas but frequently travel to rural areas to hunt, while 26 (21%) live in the rural zone. The ethical approval for the study was obtained from the Ethics committee of Hospital Lauro Wanderley (protocol number: CEP/HULW n° 103/10).

Vernacular names of the specimens cited were recorded and the animals identified in the following ways: (1) analysis of the specimens or parts thereof donated by the interviewees; (2) analysis of photographs of animals taken during the interviews and during the accompaniment of hunting activities; (3) use of identifications by taxonomists familiar with the fauna of the study area and use of vernacular names; and (4) information from previous ethnozoological studies carried out in the study area (Alves et al. 2009b; Confessor et al. 2009; Mendonça et al. 2011). The scientific nomenclature of the species that are cited in this study follows the guidelines of the Brazilian Society of Herpetology (<http://www.sbherpetologia.org.br/>).

After analysis, specimens were deposited at the zoological collections of the Universidade Federal da Paraíba. Samples were collected with the permission of the Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) and the Sistema de Autorização e Informação em Biodiversidade (SISBIO), license number 25926-2.

### Data Analysis

An accumulation curve of the herpetofaunal species cited by interviewees was prepared. In an accumulation curve for ethnobiological data, the X-axis corresponds to the number of individuals interviewed and Y-axis the number of species cited by the respondents. The curve was randomized 1,000 times and the means were calculated using the software EstimateS© version 8.2 (Colwell 2009). EstimateS© permits the statistical analysis of species richness (for this work, species richness can be interpreted as the richness of species locally exploited) of samples by determination of the Chao2 index (Colwell and Coddington 1994). This index has been used in pre-




Fig. 1. Location of the municipality of Pocinhos (Paraíba State, Northeast Brazil), where the study was conducted.

vious ethnozoological studies (Ferreira et al. 2012; Souto et al. 2011; Whiting et al. 2011).

The non-parametric estimator Chao2 (Chao 1987) is calculated by the following formula:

$$\text{Chao2} = \text{Sobs} + \left( \frac{L^2}{2M} \right)$$

where: Sobs corresponds to the number of species in a given sample, L is the number of species in only one sample (“uniques”), and M is the number of species that occur exactly in two samples. The utilization of the Chao2 estimator is recommended for ethnozoological studies since it is a non-parametric estimator based on data of incidence.

The data were entered in EstimateS© using a spreadsheet of type of respondent (rows) x type of species (columns). In preparing the spreadsheet, a value of 1 was given for each species mentioned by an interviewee and 0 for those that were not recorded.

For each species we calculated the Use-Value [adapted from the proposal of Phillips et al. (1994)], a quantitative method that demonstrates the relative importance of species known locally. This value was calculated using the following formula:  $UV = \sum U/n$ , where: UV = Use-Value of the species; U = number of citations per species; n = number of informants. The calculations of the Use-Values of any species is based objectively on the importance attributed by the informants themselves, and does not depend on the opinion of the researcher.

## Results

We recorded 18 species of herpetofauna (17 reptiles and 1 amphibian) that interacted with people in the surveyed area, either because they have some utilitarian value or because they are involved in conflicting relations with local inhabitants (Table 1). Products derived from herpetofauna were used for the following purposes: food ( $n = 7$  species), medicinal use ( $n = 7$  species), pets ( $n = 4$  species), ornamental use ( $n = 4$  species), and commerce ( $n = 2$  species). Additionally, 13 species were hunted because they are considered harmful (particularly snakes), although some of these also provide products of utilitarian value.


Based on the data collected, the mean number of species observed (Sobs) was compared with that expected to be cited in the surveyed area (Fig. 2). The results demonstrated that the sampling efficiency was adequate, since 78.4% of all species of the herpetofauna of ethnozoological importance for the study area ( $\text{Chao2} = 22.96 \pm 5.07$ ) were recorded. The species accumulation curve showed a tendency to stabilize.

When we considered the utilitarian value of the herpetofauna in the area studied, a greater number of species

were cited for their utilization as food ( $n = 7$  species), where lizards were the principal group cited for this purpose, mainly the White tegu (*Salvator merianae*, Dumeril and Bibron 1839; Use-Value = 0.66). Other lizards reported as being used for food were the Green iguana (*Iguana iguana*, Linnaeus 1758) and the whiptail lizard (*Ameivula ocellifera*, Spix 1825), with the latter being rarely consumed, as it was cited by only two interviewees. In relation to snakes, only three hunters cited species useful as food: rattlesnake (*Crotalus durissus*, Linnaeus 1758) and Rainbow boa (*Epicrates assisi*, Machado 1945). The Northeastern pepper frog (*Leptodactylus vastus*, Lutz 1930) is the only amphibian used for food according to interviewees.

The medicinal use of herpetofauna, reported by 28 hunters, appears to be the most common form of utilization for this animal group. The species most utilized for this purpose, according to the interviewees, are the White tegu ( $n = 28$  citations), Green iguana ( $n = 14$  citations), and rattlesnake ( $n = 8$  citations; Table 2). From the animals cited as useful in popular medicine, various parts or medicinal subproducts are extracted, especially the fat and hide, which are used in the treatment of various diseases and are administered in various ways (Table 2).

Use of reptiles as pets was recorded in only three of the homes visited, suggesting that the use of herpetofauna as pets is not a common practice in the study area. Species used as pets were: Red footed tortoise (*Chelonoidis carbonaria*, Spix 1824; raised by three hunters), Tuberculate toadhead turtle (*Mesoclemmys tuberculata*, Lüderwaldt 1926), White tegu, and Boa snake (*Boa constrictor*, Linnaeus 1758; cited by only one hunter). The hunter who mentioned this last species stated that he captured the animal by hand on a hunting trip, but that he did not keep the animal long at his home because he was unable to feed it adequately, thus letting it go in the forest.



**Fig. 2.** Graphs showing the values obtained with the richness estimators of herpetofaunal species hunted in surveyed area. Number of Species Observed (Sobs =  $18 \pm 2.44$ ), Number of species estimated (Chao2 =  $22.96 \pm 5.07$ ).

**Table 1.** Hunted herpetofaunal species with their respective popular names in the surveyed area. Legend: **F** = food resource, **M** = medicinal, **C** = commerce, **P** = pets, **O** = ornamentation and decoration, and **CR** = conflicting relationships.

Family/species/popular name	Use-Value	Uses and/or conflicting relationships					
		F	M	C	P	O	CR
<b>Leptodactylidae</b>							
<i>Leptodactylus vastus</i> (Lutz, 1930) – “Jia,” Northeastern pepper frog	0.01	X	X				
<b>Testudinidae</b>							
<i>Chelonoidis carbonaria</i> (Spix, 1824) – “Jabuti,” Red footed tortoise	0.01		X		X		
<b>Chelidae</b>							
<i>Mesoclemmys tuberculata</i> (Luederwaldt, 1926) – “Cágado d’água,” Tuberculate toadhead turtle	0.008		X	X			
<b>Iguanidae</b>							
<i>Iguana iguana</i> (Linnaeus, 1758) – “Camaleão,” Common green iguana	0.20	X	X	X			
<b>Teiidae</b>							
<i>Ameivula ocellifera</i> (Spix, 1825) – “Calango,” Spix’s whiptail	0.008	X					
<i>Salvator merianae</i> (Duméril and Bibron, 1839) – “Teju,” White tegu	0.66	X	X	X	X	X	X
<b>Boidae</b>							
<i>Boa constrictor</i> (Linnaeus, 1758) – “Cobra de veado,” “jibóia,” Boa snake	0.03	X	X			X	X
<i>Epicrates assisi</i> (Linnaeus, 1758) – “Salamanta,” Rainbow boa	0.16	X				X	X
<b>Colubridae</b>							
<i>Oxybelis aeneus</i> (Wagler, 1824) – “Cobra de cipó,” Brown vine snake	0.008						X
<b>Dipsadidae</b>							
<i>Boiruna sertaneja</i> (Zaher, 1996) – “Cobra preta,” Black snake	0.02						X
<i>Philodryas olfersii</i> (Lichtstein, 1823) – “Cobra verde,” Lichtenstein’s Green racer	0.02						X
<i>Leptodeira annulata</i> (Linnaeus, 1758) – “Jararaca,” Salamanta de parede, Banded cat-eyed snake	0.11						X
<i>Philodryas nattereri</i> (Steindachner, 1870) – “Cobra corre campo,” Paraguay green racer	0.04						X
<i>Pseudoboa nigra</i> (Duméril, Bibron e Duméril, 1854) – “Cobra de leite,” Black false boa	0.01						X
<i>Xenodon merremii</i> (Wagler, 1824) – “Jararaquinha,” “Goipeba,” Wagler’s snake	0.01						X
<b>Elapidae</b>							
<i>Micrurus ibiboboca</i> (Merrem, 1820) – “Cobra coral,” Caatinga coral snake	0.11						X
<b>Viperidae</b>							
<i>Bothrops erythromelas</i> (Amaral, 1923) – “Malha de cascavel,” Jararaca da seca, Caatinga lancehead	0.02						X
<i>Crotalus durissus</i> (Linnaeus, 1758) – “Cascavel,” South American rattlesnake	0.20	X	X	X		X	X

The use of herpetofauna to make artisanal products was mentioned by only three interviewees, where the hide is the principal product used for this purpose. This product is used mainly in the manufacture of accessories (belts, purses, and key chains). The species used for this purpose are: rattlesnake, whose rattle is used in the manufacture of key chains by some hunters and the hide, which can be used to make belts; and Boa snake, Rainbow boa, and White tegu, whose hide is used in the manufacture of accessories.

Despite being sources of products used for different purposes, the main motivation for the hunting and killing of the herpetofauna in the study region is that many of the species cited are considered harmful, particularly the snakes, considered venomous and efficient predators that

pose a risk to humans and their domestic animals. Forty (32.2%) hunters interviewed affirmed having killed some type of reptile while hunting or during daily activities in the countryside. Meanwhile, the hunters were unanimous in stating that they kill whatever snake they encounter. The most persecuted species are the rattlesnake ( $n = 26$  citations), Rainbow boa ( $n = 21$  citations), Caatinga lancehead (*Bothrops erythromelas*, Amaral 1923;  $n = 3$  citations), and coral snake (*Micrurus ibiboboca*, Merrem 1820;  $n = 14$  citations).

Besides snakes, the White tegu can be killed by some hunters ( $n = 4$ ) of rural areas because they do damage, since this lizard feeds on chicks and chicken eggs. The latter are important food for local families, besides being a source of income when sold.

**Table 2.** Herpetofauna used for medicinal purposes cited by hunters in the Pocinhos city, Paraíba State, Brazil.

Species / vernacular name	Citations	Medicinal use (Treated diseases)	Parts
<i>Chelonoidis carbonaria</i> (Spix, 1824)	2	Rheumatism and swelling	Shell and fat
<i>Mesoclemmys tuberculata</i> (Luederwaldt, 1926)	2	Sore throat, cough, asthma, earache, wounds, rheumatism, haemorrhoids, shortness of breath, bronchitis	Fat
<i>Iguana iguana</i> (Linnaeus, 1758)	14	Suck a splinter out of skin or flesh, snakebite, choking, boils, rheumatism, earache, sore throat, and wounds	Skin, fat, and bone
<i>Salvator merianae</i> (Duméril e Bibron, 1839)	28	Sore throat, earache, choking, deafness, boils, wounds, arthritis, asthma, rheumatism, headache, tumor; suck a splinter out of skin or flesh, cough, and swelling	Fat, tongue, and skin
<i>Crotalus durissus</i> (Linnaeus, 1758)	8	Asthma, sore throat, skin problems, cancer, rheumatism, urinary problems, arthritis, toothache, haemorrhoids, backache, mycoses, wounds, deafness, and varicose veins	Rattle and fat
<i>Leptodactylus vastus</i> (Lutz, 1930)	1	Sore throat	Meat

## Discussion

Our results reveal that the people of the surveyed area establish a greater interaction with reptiles than amphibians. This finding can be related to the greater richness of reptiles that occurs in the Caatinga (140 reptiles and 65 amphibians) and also among the reptiles there are larger-sized species, which can offer larger amounts of products for use. Snakes are feared animals in all of the semiarid northeast and in other places in Brazil, calling extra attention associated with the prevention of potential accidents (Alves et al. 2010b, 2012b, c; Moura et al. 2010).

Despite the negative view related to the many species of reptiles in the area studied, there are many species (even those killed because of conflicts) that supply products used by the local inhabitants. These observations are in agreement with Marques (1995), who noted that the link between humans and animals is fraught with contradictions and ambiguities, as the native fauna can represent either a resource or a risk to the local people.

The small number of species of herpetofauna used as food is not surprising, since traditionally, this group does

not play an important role as a protein source for the populations living in the Caatinga. The principal groups of wild vertebrates used as a source of protein in the region are birds and mammals (Alves et al. 2009b; Bezerra et al. 2011, 2012a, b, 2013; Fernandes-Ferreira et al. 2012b). However, the game importance of the White tegu should be pointed out, as its meat is used as a source of protein in the Caatinga. Such observation can be substantiated in a parallel study on the consumption of bushmeat in the same area as the present study (Mendonça 2012), which monitored the consumption of meat by local families during a year and within the local herpetofauna, only recording the consumption of two species of reptiles: *S. merianae* and *I. iguana*, with greater frequency for the former. The use of these two species for food also has been recorded in other localities of the semiarid northeast, including urban areas (Alves et al. 2012a; Marques and Guerreiro 2007). Considering the cultural and utilitarian importance of the lizard *S. merianae*, we do not exaggerate when we suggest that this animal represents one of the animals of greatest ethnozoological importance in the Brazilian Caatinga. This can be due to its size, since it is the largest species of lizard of the semiarid region (Vanzolini et al. 1980) and corroborates the findings of Alves et al. (2012b), which pointed out that *S. merianae* represents the main game reptile of the semi-arid region of Brazil.

Corroborating a tendency observed in other studies (Alves et al. 2012c; Marques and Guerreiro 2007; Santos-Fita et al. 2010), the consumption of snakes was little cited by the hunters in the study area. In Brazil, only five snake species have been reported as being used for human consumption: *Boa constrictor*, *Eunectes murinus*, *Lachesis muta*, *Crotalus durissus*, and *Epicrates asisi* (Alves et al. 2012c; Fernandes-Ferreira et al. 2013). Alves et al. (2012c) highlighted that the small numbers of snake species currently used as food in Brazil is not surprising given the negative images attributed to these animals in myths, legends, and popular beliefs. Reinforcing this notion, Rea (1981) noted that not only are snakes rejected because of their disagreeable nature but also any other creature with a similar shape or behavior. A study undertaken among human populations living along the banks of the Rio Negro (Amazonas State, Brazil) indicated that the electric eel (*Electrophorus electricus*) was one of the least favored meats because of its strong smell and the shape of its body—“it looks just like a snake” (Silva 2007).

Although the herpetofauna does not play an important role as a source of protein in the region studied, this group stands out when considering the popular medicine of the region. Despite having been cited less as medicinal species ( $n = 6$  species) than as those used for food ( $n = 7$  species), medicinal use showed a higher number of citations, suggesting its greater dissemination among the interviewees. In this context, the tegu was also featured with regard to number of citations as well its broad



medicinal applicability. Studies in various localities have already indicated the importance of this species of lizard in popular medicine in Brazil (Alves 2009; Alves et al. 2007, 2009a, 2011; Ferreira et al. 2012; Oliveira et al. 2010), even in urban areas, where the sale of products derived from *S. merianae* (as well as other species of reptiles recorded in this work) is common in public markets in various cities in northeast Brazil (Alves and Rosa 2007; Alves and Rosa 2010; Ferreira et al. 2012).

Raising wild animals as pets, particularly wild birds (Alves et al. 2010a, 2013a; Bezerra et al. 2001, 2013; Fernandes-Ferreira et al. 2012b; Nobrega et al. 2012) is a very common practice in the semiarid northeastern region, but few species of the herpetofauna are utilized for this reason, in accordance with our finding presented here. Among the reptiles of the Caatinga, the Jabuti (*C. carbonaria*) is one of the species of the most popular pets, probably because it is considered docile and easy to capture and keep in captivity. Additionally, there is also a popular belief that its presence helps avoid illnesses such as bronchitis and asthma (Alves et al. 2009a).

The strong aversion to reptiles, especially snakes, is common in various places in Brazil (Alves et al. 2012b, c; Moura et al. 2010; Santos-Fita et al. 2010), and was also recorded in our study. This aversion serves as a strong motivation for hunters and the public in general to kill snakes indiscriminately, where they are persecuted and killed whenever they are encountered. People are used to killing not only venomous snakes but also the non-venomous species, and even those amphibians that have a similar body shape as snakes. Similarly, Santos-Fita et al. (2010) documented that all inhabitants of a semiarid area of the state of Bahia have strong negative reactions in relation to snakes, always killing them if possible. It should be emphasized that these conflicts involve other groups besides snakes. In our study, we recorded that even species of reptiles with high utilitarian value, such as the tegu, can also be killed for feeding on chicken eggs, causing financial losses for farmers.

Our data, together with previous findings of other ethnozoological studies carried out in the semiarid region of the northeast, allow us to suggest some patterns of interactions between the people and herpetofauna of the Caatinga: (1) there are more frequent interactions between the people and reptiles than with amphibians; (2) lizards comprise the group with the most important species for food, particularly the White tegu; (3) products from herpetofauna play an important role in popular medicine in the semiarid, northeastern region; (4) besides food and medicinal use, products from herpetofauna can be used in handicrafts and jewelry; and, (5) various reptile species, especially snakes, are hunted and killed because of cultural aversion to these animals and the risks they pose to people and domestic animals.

Information from previous studies and that obtained here demonstrate that in the semiarid region of Brazil's northeast, reptiles and amphibians are hunted because

they are useful or considered dangerous, and sometimes for both reasons. The implementation of conservation measures aimed at the herpetofauna in this region is particularly difficult due to the aversion of the people to a good part of the species of this group. Therefore, strategies of environmental education should be adopted, besides specific actions directed at species of high game value, taking into consideration the cultural, social, and utilitarian role that governs the interactions of human populations and the herpetofauna of the Caatinga.

## Literature Cited

- Albuquerque UP, Araújo EL, Souto A, Bezerra B, Freire EMX, Sampaio E, Casas FL, Moura G, Pereira G, Melo JG, Alves M, Rodal M, Schiel N, Neves RL, Alves RRN, Azevedo-Júnior S, Telino Júnior W. 2012. Caatinga revisited: Ecology and conservation of an important seasonal dry forest. *Scientific World Journal* 2012(205182): 1–18.
- Alves RRN. 2009. Fauna used in popular medicine in Northeast Brazil. *Journal of Ethnobiology and Ethnomedicine* 5(1): 1–30. doi:10.1186/1746-4269-5-1
- Alves RRN. 2012. Relationships between fauna and people and the role of ethnozoology in animal conservation. *Ethnobiology and Conservation* 1: 1–69
- Alves RRN, Barbosa JAA, Santos SLDX, Souto WMS, Barboza RRD. 2011. Animal-based remedies as complementary medicines in the semi-arid region of northeastern Brazil. *Evidence-based Complementary and Alternative Medicine* 2011(179876): 1–15. doi:10.1093/ecam/nep134
- Alves RRN, Gonçalves MBR, Vieira WLS. 2012a. Caça, uso e conservação de vertebrados no semiárido Brasileiro. *Tropical Conservation Science* 5(3): 394–416.
- Alves RRN, Léo Neto NA, Santana GG, Vieira WLS, Almeida WO. 2009a. Reptiles used for medicinal and magic religious purposes in Brazil. *Applied Herpetology* 6(3): 257–274.
- Alves RRN, Lima JRF, Araújo HF. 2013a. The live bird trade in Brazil and its conservation implications: An overview. *Bird Conservation International* 23 (01): 53–65. doi:10.1017/S095927091200010X
- Alves RRN, Mendonça LET, Confessor MVA, Vieira WLS, Lopez LCS. 2009b. Hunting strategies used in the semi-arid region of northeastern Brazil. *Journal of Ethnobiology and Ethnomedicine* 5 (12):1–50. doi:10.1186/1746-4269-5-12
- Alves RRN, Nogueira E, Araújo H, Brooks S. 2010a. Bird-keeping in the Caatinga, NE Brazil. *Human Ecology* 38(1): 147–156
- Alves RRN, Pereira-Filho GA, Vieira KS, Santana GG, Vieira WLS, Almeida WO. 2010b. Répteis e as populações humanas no Brasil: uma abordagem etnoherpetológica. Pp. 121–146 In: Editors, Alves RRN, Souto WMS, Mourão JS. A Etnozootologia no Brasil:

- Importância, status atual e perspectivas futuras Ed. NUPEEA, Recife, Brazil.
- Alves RRN, Pereira Filho GA, Vieira KS, Souto WMS, Mendonça LET, Montenegro PFGP, Almeida WO, Vieira WLS. 2012b. A zoological catalogue of hunted reptiles in the semiarid region of Brazil. *Journal of Ethnobiology and Ethnomedicine* 8(17): 1-29. doi:10.1186/1746-4269-8-27.
- Alves RRN, Rosa IL. 2007. Zotherapy goes to town: The use of animal-based remedies in urban areas of NE and N Brazil. *Journal of Ethnopharmacology* 113: 541–555. doi:10.1016/j.jep.2007.07.015
- Alves RRN, Rosa IL. 2010. Trade of animals used in Brazilian traditional medicine: trends and implications for conservation. *Human Ecology* 38(5): 691–704.
- Alves RRN, Rosa IL, Santana GG. 2007. The role of animal-derived remedies as complementary medicine in Brazil. *BioScience* 57(11): 949–955.
- Alves RRN, Souto WMS. 2011. Ethnozology in Brazil: Current status and perspectives. *Journal of Ethnobiology and Ethnomedicine* 7(22): 1-18. doi:10.1186/1746-4269-7-22
- Alves RRN, Vieira KS, Santana GG, Vieira WLS, Almeida WO, Souto WMS, Montenegro PFGP, Pezzuti JCB. 2012c. A review on human attitudes towards reptiles in Brazil. *Environmental Monitoring and Assessment* 184(11): 6877–6901. doi: 10.1007/s10661-011-2465-0.
- Alves RRN, Vieira WLS, Santana GG. 2008. Reptiles used in traditional folk medicine: Conservation implications. *Biodiversity and Conservation* 17: 2037–2049. doi:10.1007/s10531-007-9305-0.
- Alves RRN, Vieira WLS, Santana GG, Vieira KS, Montenegro PFGP. 2013b. Herpetofauna used in traditional folk medicine: Conservation Implications. Pp. 109–133 In: Editors, Alves RRN, Rosa IL. *Animals in Traditional Folk Medicine: Implications for conservation*. Volume 1. Springer-Verlag, Berlin Heidelberg, Germany.
- Bailey K. 1994. *Methods of Social Research*. Free Press, New York, New York, USA.
- Barbosa JAA, Nobrega VA, Alves RRN. 2011. Hunting practices in the semiarid region of Brazil. *Indian Journal of Traditional Knowledge* 10(3): 486-490.
- Bernard R. 2006. *Research Methods in Anthropology: Qualitative and quantitative approaches*. Ed. Altamira Press, Oxford, UK.
- Bertrand H. 1997. Contribution à l'étude de l'herpétologie et de l'ethnoherpétologie en Anjou [A study on the herpetology and ethnoherpetology of Anjou province (France)]. *Bulletin de la Société herpétologique de France* (82-83): 51–62.
- Bezerra DMM, Araújo HFP, Alves ÂGC, Alves RRN. 2013. Birds and people in semiarid northeastern Brazil: Symbolic and medicinal relationships. *Journal of Ethnobiology and Ethnomedicine* 9(3): 1-11. doi:10.1186/1746-4269-9-3
- Bezerra DMM, Araújo HFP, Alves RRN. 2012a. Captura de aves silvestres no semiárido brasileiro: técnicas cinegéticas e implicações para conservação. *Tropical Conservation Science* 5(1): 50–66.
- Bezerra DMM, Araújo HFP, Alves RRN. 2012b. Wild birds as source of food in the semi-arid region of Rio Grande do Norte State, Brazil. *Sitientibus Série Ciências Biológicas* 11(2): 177–183.
- Bezerra DMMSQ, Araújo HFP, Alves RRN. 2011. The use of wild birds by rural communities in the semi-arid region of Rio Grande do Norte State, Brazil. *Bio-remediation, Biodiversity and Bioavailability* 5(1): 117–120.
- Chao A. 1987. Estimating the population size for capture-recapture data with unequal catchability. *Biometrics* 43(4): 783–791.
- Colwell RK. 2009. EstimateS: Statistical estimation of species richness and shared species from samples. Version 8.2. User's Guide and application published at: <http://purl.oclc.org/estimates>. Storrs, Connecticut, USA.
- Colwell RK, Coddington JA. 1994. Estimating terrestrial biodiversity through extrapolation. *Philosophical Transactions of the Royal Society B* 345: 101–118.
- Confessor MVA, Mendonca LET, Mourao JS, Alves RRN. 2009. Animals to heal animals: Ethnoveterinary practices in semi-arid region, Northeastern Brazil. *Journal of Ethnobiology and Ethnomedicine* 5(37): 1-9. doi:10.1186/1746-4269-5-37.
- Das I. 1998. *The Serpent's Tongue: A contribution to the ethnoherpetology of India and adjacent countries*. Ed. Chimaira, Cornell, USA.
- Fernandes-Ferreira H, Cruz RL, Borges-Nojosa DM, Alves RRN. 2012a. Folklore concerning snakes in the Ceará State, northeastern Brazil. *Sitientibus Série Ciências Biológicas* 11(2): 153–163.
- Fernandes-Ferreira H, Mendonça SV, Albano C, Ferreira FS, Alves RRN. 2012b. Hunting, use and conservation of birds in Northeast Brazil. *Biodiversity and Conservation* 21: 221–244. doi: 10.1007/s10531-011-0179-9
- Fernandes-Ferreira H, Mendonca SV, Cruz RL, Borges-Nojosa DM, Alves RRN. 2013. Hunting of herpetofauna in montane, coastal, and dryland areas of Northeastern Brazil. *Herpetological Conservation and Biology* 8: 652–666.
- Ferreira FS, Albuquerque UP, Coutinho HDM, Almeida WO, Alves RRN. 2012. The Trade in Medicinal Animals in Northeastern Brazil. *Evidence-based Complementary and Alternative Medicine* 2012: 1–20. doi:10.1155/2012/126938
- Franke J, Telecky TM. 2001. *Reptiles as Pets: An examination of the trade in live reptiles in the United States*. Humane Society of the United States, Washington, DC, USA.
- Goodman SM, Hobbs J. 1994. The distribution and ethnozology of reptiles of the northern portion of the Egyptian eastern desert. *Journal of Ethnobiology* 14:

- 75–100.
- Huntington HP. 2000. Using traditional ecological knowledge in science: Methods and applications. *Ecological Applications* 10(5): 1270–1274.
- Klemens MW, Thorbjarnarson JB. 1995. Reptiles as a food resource. *Biodiversity and Conservation* 4(3): 281–298.
- Marques JGW. 1995. *Pescando pescadores: etnoecologia abrangente no baixo São Francisco alagoano*. NUPAUB-USP, São Paulo, Brazil.
- Marques JGW, Guerreiro W. 2007. Répteis em uma Feira Nordestina (Feira de Santana, Bahia). Contextualização Progressiva e Análise Conexivo-Tipológica. *Sitientibus Série Ciências Biológicas* 7(3): 289–295.
- Mendonça LET. 2012. Atividades cinegéticas por populações rurais e urbanas do município de Pocinhos, Estado da Paraíba, Brasil: uma abordagem etnozoológica. Universidade Federal da Paraíba, João Pessoa, Brasil.
- Mendonça LET, Souto CM, Andreilino LL, Souto WMS, Vieira WLS, Alves RRN. 2011. Conflitos entre pessoas e animais silvestres no semiárido paraibano e suas implicações para conservação. *Sitientibus Série Ciências Biológicas* 11(2): 185–199.
- Morris R, Morris D. 1965. *Men and Snakes*. McGraw-Hill, New York, New York, USA.
- Moura MR, Costa HC, São-Pedro VA, Fernandes VD, Feio RN. 2010. The relationship between people and snakes in eastern Minas Gerais, southeastern Brazil. *Biota Neotropica* 10(4): 133–141.
- Nobrega VA, Barbosa JAA, Alves RRN. 2012. Use of wild birds by residents of the municipality of Fagundes, in Paraíba's semiarid region: An ethnoornitologic approach. *Sitientibus Série Ciências Biológicas* 11(2): 165–175.
- Oliveira ES, Torres DF, Brooks SE, Alves RRN. 2010. The medicinal animal markets in the metropolitan region of Natal City, Northeastern Brazil. *Journal of Ethnopharmacology* 130(1): 54–60.
- Phillips O, Gentry AH, Reynel C, Wilkin P, Galvez-Durand BC. 1994. Quantitative ethnobotany and Amazonian conservation. *Conservation Biology* 8: 225–248.
- Rea AM. 1981. Resource utilization and food taboos of Sonoran desert peoples. *Journal of Ethnobiology* 1: 69–83.
- Ribeiro RS. 2003. *Pocinhos: o local e o geral*. RG Gráfica e Editora, Campina Grande, Brasil.
- Santos-Fita D, Costa-Neto EM, Schiavetti A. 2010. 'Offensive' snakes: Cultural beliefs and practices related to snakebites in a Brazilian rural settlement. *Journal of Ethnobiology and Ethnomedicine* 6(1): 1–13. doi:10.1186/1746-4269-6-13
- Schlaepfer MA, Hoover C, Dodd Jr CK. 2005. Challenges in evaluating the impact of the trade in amphibians and reptiles on wild populations. *BioScience* 55(3): 256–264.
- Silva AL. 2007. Comida de gente: Preferências e tabus alimentares entre os ribeirinhos do Médio Rio Negro (Amazonas, Brasil). *Revista de Antropologia* 50: 125–179.
- Souto WMS, Mourão JS, Barboza RRD, Mendonça LET, Lucena RFP, Confessor MVA, Vieira WLS, Montenegro PFGP, Lopez LCS, Alves RRN. 2011. Medicinal animals used in ethnoveterinary practices of the 'Cariri Paraibano', NE Brazil. *Journal of Ethnobiology and Ethnomedicine* 7(30): 1–19. doi:10.1186/1746-4269-7-30.
- Speck FG. 1946. Ethnoherpetology of the Catawba and Cherokee Indians. *Journal of the Washington Academy of Sciences* 36: 355–360.
- Vanzolini PE, Ramos-Costa AMM, Vitt LJ. 1980. Répteis das caatingas. Academia Brasileira de Ciências, Rio de Janeiro, Brazil.
- Whiting MJ, Williams VL, Hibbitts TJ. 2011. Animals traded for traditional medicine at the Faraday market in South Africa: species diversity and conservation implications. *Journal of Zoology* 284: 84–96.

Received: 17 January 2014

Accepted: 20 June 2014

Published: 05 July 2014



## Ethnoherpetology of the Caatinga region of Brazil



**Livia Emanuelle Tavares Mendonça** is a biologist, M.Sc. in Zoology at the Universidade Federal da Paraíba (UFPB), Bachelor in Biological Sciences from the Universidade Estadual da Paraíba (UEPB). She develops research focused on hunting, wildlife conservation, and Ethnozoology (Photographed by Rômulo Alves).



**Washington Luiz Silva Vieira** is a biologist, Ph.D. in Zoology at the Universidade Federal da Paraíba. His main interests are taxonomy, ecology, and natural history of herpetofauna. He also has research interests in ethnoherpetology and animal conservation (Photographed by Rômulo Alves).



**Rômulo Romeu Nóbrega Alves** is a professor at the Universidade Estadual da Paraíba, Brazil, where he teaches undergraduate and graduate courses in Biological Sciences and Ecology. His Ph.D. (Zoology) was completed in 2006, at the Universidade Federal da Paraíba. His areas of professional interest are ethnozoology and wildlife trade, uses and conservation, zotherapy, and human ecology. He has conducted ethnobiological research for the last ten years in Brazil which focuses on fisheries, hunting, and wildlife trade and uses. Currently, he coordinates projects on hunting and uses of wildlife in Brazil. In addition, he is one of the Editors-in-Chief of the journal of *Ethnobiology and Conservation* and of the Editorial Board of the *Journal of Ethnobiology and Ethnomedicine*. Prof. Rômulo R. N. Alves holds a Productivity scholarship, provided by the National Council of Science and Technology (CNPq) (Photographed by Wedson Souto).